



Dung beetles of Azorean cattle-grazed pasturelands - data of the DUNGPOOL project

Almudena Duenas-Rojas^{‡,§}, Eva Cuesta^{‡,§}, Laurine M. Parmentier[|], Abrão Leite[¶],
Paulo A. V. Borges^{|,#,□}, Ana M. C. Santos^{‡,§,|}

‡ Centro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Universidad Autónoma de Madrid, Calle Darwin 2, 28049, Madrid, Spain

§ Terrestrial Ecology Group (TEG-UAM), Departamento de Ecología, Universidad Autónoma de Madrid, Calle Darwin 2, Edificio de Biología, 28049, Madrid, Spain

| University of the Azores, cE3c- Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity Group, CHANGE – Global Change and Sustainability Institute, School of Agricultural and Environmental Sciences, Rua Capitão João d'Ávila, Pico da Urze, 9700-042, Angra do Heroísmo, Azores, Portugal

¶ Rua Fernando Pessoa, n. 99 2765-483, Estoril, Portugal

IUCN SSC Atlantic Islands Invertebrate Specialist Group, 9700-042, Angra do Heroísmo, Azores, Portugal

□ IUCN SSC Species Monitoring Specialist Group, 9700-042, Angra do Heroísmo, Azores, Portugal

Corresponding author: Almudena Duenas-Rojas (almudena.duenas@uam.es)

Academic editor: António O. Soares

Received: 26 Jun 2025 | Accepted: 19 Sep 2025 | Published: 01 Oct 2025

Citation: Duenas-Rojas A, Cuesta E, Parmentier LM, Leite A, Borges PAV, Santos AMC (2025) Dung beetles of Azorean cattle-grazed pasturelands - data of the DUNGPOOL project. Biodiversity Data Journal 13: e163289.

<https://doi.org/10.3897/BDJ.13.e163289>

Abstract

Background

The data we present are part of the project DUNGPOOL, which aims to evaluate the effects of the species pool, community assembly processes and increasing temperatures on the local diversity and ecosystem functions performed by dung beetles in island and mainland cattle-grazed pasturelands. By combining replicated field experiments on the Iberian Peninsula with parallel work on three mid-Atlantic islands (Pico, Faial and Terceira, Azores), the project exploits the natural contrast between a species-rich mainland pool and the species-poor, largely exotic island pools, to test explicit biodiversity–ecosystem–function (BEF) hypotheses across spatial scales.

From June to July 2024, dung beetles were sampled in 84 locations of the three islands of the Azores Archipelago (Portugal), using 306 pitfall traps baited with fresh cow dung.

New information

A total of 1,141 dung beetle individuals were recorded across these islands: 18 individuals in Pico Island, 480 individuals in Faial Island and 643 individuals in Terceira Island. These individuals were distributed amongst four species from the family Scarabaeidae (Insecta, Coleoptera): *Calamosternus granarius* (Linnaeus, 1767), *Labarrus lividus* (Olivier, 1789), *Onthophagus taurus* (Schreber, 1759) and *Onthophagus medius* (Kugelann, 1792). The species *O. medius* is a new record for this Archipelago and we propose that previous historical records of *Onthophagus vacca* (Linnaeus, 1767) should now be considered a regional synonym of *Onthophagus medius*.

Keywords

Azores Islands, Faial Island, Pico Island, Terceira Island, Coleoptera, Scarabaeidae, dung beetles, occurrence event, pastures, pitfall traps, sampling event

Introduction

Dung beetles (Coleoptera, Scarabaeoidea) are known for their coprophagous habits. They are classified into three functional groups based on their behavioural strategies: telecoprids, species that roll dung balls and bury them at a distance from the main dung pile; paracoprids, those that make galleries in the soil and bury the dung underneath the main dung pile, in which they bury small pieces of dung; and endocoprids, those that feed in the dung. As a result of these habits, dung beetles perform different ecological functions that provide important ecosystem services (Nichols et al. 2008), like the decomposition and burial of dung and secondary dispersal of seeds (deCastro-Arrazola et al. 2020, Kadiri et al. 2022, deCastro-Arrazola et al. 2023), amongst others. Human activities associated with landscape transformation and habitat fragmentation are a serious threat to dung beetles (Noriega et al. 2021, Lumaret et al. 2022). The intensification of livestock grazing alters the natural conditions of the soil and, therefore, their habitat and diversity (Noriega et al. 2023). On the other hand, the use of antiparasitics in livestock, such as ivermectins, also affects the diversity and ecosystem functions of dung beetles (Verdú et al. 2018, Ambrožová et al. 2021).

Many dung beetles are associated with cattle-grazed pasturelands, a land use of high cultural and economic importance in the Azorean Islands, Portugal (Borges et al. 2020), an isolated archipelago of volcanic origin that is located in the North Atlantic and is part of the Macaronesia Region. The Azores were discovered in the 15th century by the Portuguese and, since human colonisation, native habitats have been widely modified (Borges et al. 2020).

Humans introduced livestock in these islands and because of its expansion, many native habitats were replaced by pastures for cattle grazing. The dung beetles present in this Archipelago are mostly introduced species (Borges et al. 2022) and their presence might be very beneficial in these novel ecosystems, as these insects enhance the decomposition of dung, which can quickly accumulate in the pasturelands associated with cattle production.

Here, we present the diversity of dung beetles (Coleoptera, Scarabaeidae) associated with pasturelands in three Azorean islands (Faial, Pico and Terceira), collected within the project DUNGPOOL, which aims to evaluate the effects of the species pool, community assembly processes and increasing temperatures on the local diversity and ecosystem functions performed by dung beetles in island and mainland cattle-grazed pasturelands.

General description

Purpose: The main objective of this study is to provide a recent inventory of dung beetles species present in the pasturelands of Pico, Faial and Terceira Islands (Azores, Portugal).

Additional information: The DUNGPOOL project was conceived to unravel how the size and composition of the regional species pool, the mechanisms of community assembly (including priority effects and biotic interactions) and near-term climate warming jointly shape dung beetle diversity and the key ecosystem functions they provide in pasturelands. By combining replicated field experiments on the Iberian mainland with parallel work on three mid-Atlantic islands (Pico, Faial and Terceira, Azores), the project exploits the natural contrast between a species-rich mainland pool and the species-poor, largely exotic island pools to test explicit biodiversity–ecosystem-function (BEF) hypotheses across spatial scales.

Project description

Title: Effects of species pool and community assembly processes on dung beetle diversity and ecosystem functions in a warming world (DUNGPOOL)

Personnel:

- Principal investigator: Ana M. C. Santos
- Fieldwork (site selection and experimental setting): Paulo A.V. Borges and Ana M. C. Santos.
- Fieldwork (authorisation): Azorean Regional Directorate for the Environment (Internationally Recognized Compliance Certificate CCIR-RAA/2024/6).
- Fieldwork team:
- Pico Island: Almudena Dueñas-Rojas, Eva Cuesta, Paulo A.V. Borges, Ana M. C. Santos
- Faial Island: Almudena Dueñas-Rojas, Eva Cuesta, Laurine Parmentier, Ana M. C. Santos

- Terceira Island: Almudena Dueñas-Rojas, Eva Cuesta, Abrão Leite, Ana M. C. Santos
- Taxonomist: Almudena Dueñas-Rojas and Eva Cuesta
- Darwin Core databases: Almudena Dueñas-Rojas

Funding: Agencia Estatal de Investigación, Ministerio de Ciencia, Innovación y Universidades (Spain) (PID2021-122380NA-I00); MICIU/AEI/10.13039/501100011033; FEDER, UE.

Sampling methods

Description: This study was conducted from June to July 2024, in a total of 84 sites of three Azorean Islands: 29 in Pico Island, 29 in Faial Island and 26 in Terceira Island. Most of these sites correspond to cattle-grazed pasturelands.

Sampling description: In most sampling sites, we placed three pitfall traps filled with a soap-water solution and baited with fresh cow dung. However, in three of the sites in each island, we set nine baited pitfall traps (Fig. 1). Therefore, a total of 306 traps were placed, 105 in Pico, 105 in Faial and 96 in Terceira. The traps were collected after 72 hours of exposure.




Figure 1. [doi](#)

Sampling sites in three Azorean Islands (Portugal): Pico Island, Faial Island and Terceira Island. Blue dots correspond to sites sampled using three pitfall traps and red dots correspond to those sites in which nine pitfall traps were used.

The collected samples were stored in 70% alcohol. Then, in the laboratory they were cleaned and the dung beetles were separated. The specimens were then identified to the species level, labelled and preserved in 70% alcohol.

Quality control: All specimens have been identified by taxonomists from Universidad Autónoma de Madrid (Eva Cuesta and Almudena Dueñas-Rojas).

Geographic coverage

Description: Three islands of the Archipelago of the Azores, Portugal: Pico, Faial and Terceira (Fig. 2). Most of the area of these islands is occupied by pastures used for cattle grazing (Borges et al. 2020).



Figure 2. [doi](#)

Location and map of the Azores Islands (Portugal): Pico (38.495833, -28.455833), Faial (38.588056, -28.665833) and Terceira (decimal coordinates 38.710556, -27.160000).

Coordinates: 38.428000 and 38.793000 Latitude; -28.746000 and -27.054000 Longitude.

Taxonomic coverage

Taxa included:

Rank	Scientific Name
phylum	Arthropoda
class	Insecta

order	Coleoptera
family	Scarabaeidae
subfamily	Aphodiinae
subfamily	Scarabaeinae

Temporal coverage

Data range: 2024-6-27 - 2024-7-25.

Usage licence

Usage licence: Open Data Commons Attribution License

Data resources

Data package title: Dung beetles of Azorean cattle-grazed pasturelands – data of the DUNGPOOL project

Resource link: <https://doi.org/10.15470/ego4ut>

Alternative identifiers: <https://www.gbif.org/dataset/cec13430-b81f-4d08-b066-5fd5e523aaa8>

Number of data sets: 2

Data set name: Event Table

Character set: UTF-8

Download URL: https://ipt.gbif.es/resource?r=dungpool_azores#anchor-downloads

Data format: Darwin Core Archive format

Data format version: 1.3

Description: The dataset was published in the Global Biodiversity Information Facility platform, GBIF (Dueñas-Rojas et al. 2025). The following data table includes all the records for which a taxonomic identification of the species was possible. The dataset submitted to GBIF is structured as a sample event dataset that has been published as a Darwin Core Archive (DwCA), which is a standardised format for sharing biodiversity data as a set of one or more data tables. The core data file contains 123 records (eventID).

Column label	Column description
EventID	Identifier of the events, unique for the dataset.
locationID	Identifier of the locations, unique for the dataset.
stateProvince	The name of the next smaller administrative region than country (state, province, canton, department, region etc.) in which the Location occurs.
islandGroup	The name of the island group in which the Location occurs (Azores Archipelago).
island	The name of the island on which the Location occurs (Pico, Faial or Terceira).
country	The full, unabbreviated name of the next smaller administrative region than stateProvince (county, shire, department etc.) in which the Location occurs (Portugal).
countryCode	The standard code for the country in which the Location occurs (PT).
municipality	The full, unabbreviated name of the next smaller administrative region than county (city, municipality etc.) in which the Location occurs.
locality	The specific description of the place.
habitat	The habitat for an Event.
minimumElevationinMetres	The lower limit of the range of elevation (altitude, usually above sea level), in metres.
decimalLatitude	Approximate centre point decimal latitude of the field site in GPS coordinates.
decimalLongitude	Approximate centre point decimal longitude of the field site in GPS coordinates.
geodeticDatum	Standard Global Positioning System coordinate reference for the location of the sample collection points.
coordinateUncertaintyinMetres	Uncertain value of coordinate metrics.
coordinatePrecision	Value in decimal degrees to a precision of six decimal places.
georeferenceSources	Navigation system used to record the location of sample collections.
samplingProtocol	The sampling protocol used to capture the species.
year	Year the sample was collected.
month	The integer month in which the Event occurred.
eventDate	The date-time or interval during which an Event occurred.
sampleSizeValue	A numeric value for a measurement of the size (time duration, length, area or volume) of a sample in a sampling Event.
sampleSizeUnits	The unit of measurement of the size (time duration, length, area or volume) of a sample in a sampling Event.
eventRemarks	Comments or notes about the Event.
type	The nature or genre of the resource.

modified	The most recent date-time on which the resource was changed.
language	A language of the resource.
licence	Reference to the licence under which the record is published.
datasetID	An identifier for the set of data.
datasetName	Name of the dataset.

Data set name: Occurrence Table

Character set: UTF-8

Download URL: https://ipt.gbif.es/resource?r=dungpool_azores#anchor-downloads

Data format: Darwin Core Archive format

Data format version: 1.3

Description: The dataset was published in the Global Biodiversity Information Facility platform, GBIF (Dueñas-Rojas et al. 2025). The following data table includes all the records for which a taxonomic identification of the species was possible. The dataset submitted to GBIF is structured as an occurrence table that has been published as a Darwin Core Archive (DwCA), which is a standardised format for sharing biodiversity data as a set of one or more data tables. The core data file contains 130 records (occurrenceID).

Column label	Column description
eventID	Identifier of the events, unique for the dataset.
type	The nature or genre of the resource.
licence	Reference to the licence under which the record is published.
institutionCode	The code of the institution publishing the data.
institutionID	An identifier for the institution publishing the data.
datasetName	Name of the dataset.
basisOfRecord	The nature of the data record.
OccurrenceID	Identifier of the record, coded as a global unique identifier.
organismQuantity	A number or enumeration value for the quantity of Organisms.
organismQuantityType	The type of quantification system used for the quantity of organisms.
lifeStage	The age class or life stage of the Organism(s) at the time the Occurrence was recorded.
establishmentMeans	Statement about whether a Organism has been introduced to a given place and time through the direct or indirect activity of modern humans.

recordedBy	A list (concatenated and separated) of names of people, groups or organisations who performed the sampling in the field.
identifiedBy	A list of names of people, groups or organisations who assigned the Taxon to the subject.
dateIdentified	The date on which the subject was determined as representing the Taxon.
kingdom	Kingdom name.
phylum	Phylum name.
class	Class name.
order	Order name.
family	Family name.
subfamily	subfamily name.
scientificName	The full scientific name, with authorship and date information if known.
genus	Genus name.
specificEpithet	Specific epithet name.
scientificNameAuthorship	The authorship information for the scientificName formatted according to the conventions of the applicable nomenclaturalCode.
taxonRank	Lowest taxonomic rank of the record.
occurrenceStatus	A statement about the presence or absence of a taxon.

Additional information

A total of 1141 dung beetle individuals were recorded across three Azorean Islands (Table 1): Pico (n = 18), Faial (n = 480) and Terceira (n = 643). These individuals were distributed amongst four species: *Calamosternus granarius* (Linnaeus, 1767) (Fig. 3a), *Labarrus lividus* (Olivier, 1789) (Fig. 3b), *Onthophagus taurus* (Schreber, 1759) (Fig. 3c) and *Onthophagus medius* (Kugelann, 1792) (Fig. 3d).

Table 1.

List of species and number of individuals found on Pico Island (PIC), Faial Island (FAI) and Terceira Island (TER).

Family	Subfamily	Scientific name	PIC	FAI	TER
Scarabaeidae	Aphodiinae	<i>Calamosternus granarius</i> (Linnaeus, 1767)	1	0	1
Scarabaeidae	Aphodiinae	<i>Labarrus lividus</i> (Olivier, 1789)	0	2	33
Scarabaeidae	Scarabaeinae	<i>Onthophagus taurus</i> (Schreber, 1759)	17	478	606
Scarabaeidae	Scarabaeinae	<i>Onthophagus medius</i> (Kugelann, 1792)	0	0	3




Figure 3.

Dung beetles of Azorean cattle-grazed pastureland:

a: *Calamosternus granarius* (Linnaeus, 1767) (Credit: Javier Torrent, Azorean Biodiversity Group); [doi](#)

b: *Labarrus lividus* (Olivier, 1789) (Credit: Javier Torrent, Azorean Biodiversity Group); [doi](#)

c: *Onthophagus taurus* (Schreber, 1759) (male) (Credit: Javier Torrent, Azorean Biodiversity Group); [doi](#)

d: *Onthophagus medius* (Kugelann, 1792) (Credit: Javier Torrent, Azorean Biodiversity Group). [doi](#)

Ecological and Biogeographical implications

The survey implemented in this study revealed a very impoverished dung beetle fauna, dominated by a single species, *Onthophagus taurus*. This dominance likely reflects both the species' high ecological plasticity and superior dispersal capabilities. In a recent study, Tonelli and Lobo (2025) concluded that species with enhanced dispersal abilities and greater ecological adaptability are more prone to succeed and establish stable

populations in insular environments, where habitat diversity is often limited and ecological opportunities can be more restricted.

It is important to note, however, that the Azorean dung beetle assemblage is not limited to *O. taurus*. Other species have been recorded in the Archipelago, but were absent from our samples, such as *Onthophagus illyricus* (Scopoli, 1763) (Borges et al. 2022). The absence of *O. illyricus* in the present survey could be attributed to factors such as its lower abundance, more specific habitat requirements or possibly localised extinctions linked to habitat degradation or competition with more adaptable species like *O. taurus*. This observation aligns with broader patterns documented in island ecosystems, where a few highly adaptable species often dominate, leading to biotic homogenisation and a reduction in overall faunal diversity (Fernández-Palacios et al. 2021) and ecosystem functioning.

Moreover, after a recent revision of *O. vacca* complex, several authors concluded that a large number of past identifications of *Onthophagus vacca* (Linnaeus, 1767) belong to a different species, the very similar *Onthophagus medius* (Kugelann, 1792) (Rössner et al. 2010, Roy et al. 2016, Drăghici et al. 2019). Indeed, in Europe, two sibling species within the *Onthophagus vacca* complex can be reliably separated by a combination of information related with external morphology, seasonal timing and geographic range. Diagnostic characters are summarised in Table 1 of Rössner et al. (2010): *O. medius* typically bears larger, often confluent elytral spots and a complete basal pronotal margin, with a more truncate clypeal edge in males; in contrast, *O. vacca* has discrete, small black spots on the elytra, a pronotal basal line interrupted laterally and a sinuate male clypeal margin. Phenology further separates the two species, with adults of *O. vacca* emerging earliest in spring (activity peaking in April–May), whereas *O. medius* appears slightly later, with a narrower flight period peaking in May (Rössner et al. 2010). Biogeography also differs: *O. vacca* extends into northern Africa and is largely absent north of 50° N in central Europe, whereas *O. medius* occurs further north and at higher altitudes, but is absent from Mediterranean islands such as Corsica and Sardinia (Rössner et al. 2010). A careful re-examination of all Azorean material available in the University of Azores Insect Collection ("Dalberto Teixeira Pombo") with specimens collected by Borges (1989), Borges and Serrano (1989) and Borges et al. (2021) shows that all specimens, formerly cited as *O. vacca*, actually conform to *O. medius*. Moverover, we also examined more than 44 specimens recently collected in Terceira Island under the scope of an academic thesis (Mediavilla 2024) and the same applies. Therefore, possibly no genuine *O. vacca* have been detected in the Azores. Accordingly, we suggest that *Onthophagus medius* (Kugelann, 1792) is the sole valid member of the *O. vacca* complex in the Azorean fauna.

Notably, the general abundance of Azorean dung-beetle fauna is low and we may ask for the reasons for such pattern. In the Azores — based on 30 composite agricultural-soil samples collected between 2018 and 2020 — organic-contaminant loads are lower than in the other Macaronesian Archipelagoes, but still noteworthy in several categories (Acosta-Dacal et al. 2022). The one pharmaceutical-active compound in Azorean soils that originates from veterinary treatment of cattle is oxfendazole, a broad-spectrum anthelmintic commonly administered to cows. It was detected in 10% of the Azores

samples, indicating its use in livestock and subsequent entry into soils via manure and pasture fertilisation (Acosta-Dacal et al. 2022). Modern livestock production almost universally relies on endectocides (e.g. ivermectin, moxidectin, doramectin) to control gastrointestinal nematodes, external parasites and flies. However, nothing is known in Azores about the consequences of modern livestock on coprophagous insects.

Acknowledgements

We gratefully acknowledge the collaboration and support of all the farmers that allowed the implementation of this study (Hélio Carlos, Sebastião Dutra, José António Furtado, Paulo César Furtado, Pedro Garcia, Rui Matos, Francisco Natálio). We are also grateful to the Agriculture Services of Faial Island (Célia Mesquita, Luis Machado e Bruno Melo), Pico Island Director of Nature Park Vanda Serpa and Nature Park Rangers, Filipa Alves, Lidia Nogueira and Paulo Freitas (Serviço de Ambiente e Ação Climática do Pico / Parque Natural do Pico) and Rui Cardoso from the Terceira Island Farmers Association.

This project (PID2021-122380NA-I00) was funded by Agencia Estatal de Investigación, Ministerio de Ciencia, Innovación y Universidades (Spain), MICIU/AEI/10.13039/501100011033 and by FEDER, UE. ADR was supported by the PhD grant PRE2022-105356, funded by MCIN/AEI/10.13039/501100011033 and by FSE+. AMCS was supported by a Spanish Ramón y Cajal fellowship RYC2020-029407-I, funded by MICIU/AEI/10.13039/501100011033 and by “ESF Investing in your future”.

Author contributions

Conceptualisation: Ana M. C. Santos; Data curation: Almudena Dueñas-Rojas; Formal analysis: Almudena Dueñas-Rojas; Funding acquisition: Ana M. C. Santos; Investigation: Almudena Dueñas-Rojas, Eva Cuesta, Abrão Leite, Laurine M. Parmentier, Paulo A.V. Borges and Ana M. C. Santos; Methodology: Ana M. C. Santos; Project administration: Ana M. C. Santos and Paulo A.V. Borges; Supervision: Ana M. C. Santos and Paulo A.V. Borges; Writing – original draft: Almudena Dueñas-Rojas; Writing – review and editing: All authors.

References

- Acosta-Dacal A, Hernández-Marrero ME, Rial-Berriel C, Díaz-Díaz R, Bernal-Suárez MdM, Zumbado M, Henríquez-Hernández LA, Boada LD, Luzardo OP (2022) Comparative study of organic contaminants in agricultural soils at the archipelagos of the Macaronesia. *Environmental Pollution* 301: 118979. <https://doi.org/10.1016/j.envpol.2022.118979>
- Ambrožová L, Sládeček FXJ, Zítek T, Perlík M, Kozel P, Jirků M, Čížek L (2021) Lasting decrease in functionality and richness: Effects of ivermectin use on dung beetle communities. *Agriculture, Ecosystems & Environment* 321: 107634. <https://doi.org/10.1016/j.agee.2021.107634>

- Borges PAV (1989) Sobre alguns coleópteros (Insecta, Coleoptera) colhidos na Graciosa por Dalberto Teixeira Pombo. Relatórios e Comunicações do Departamento de Biologia 17: 57-59. URL: <http://hdl.handle.net/10400.3/864>
- Borges PAV, Serrano ARM (1989) New records of the coleopterous fauna (Insecta, Coleoptera) from the Azores. Boletim do Museu Municipal do Funchal 41: 5-24. URL: <http://hdl.handle.net/10400.3/1887>
- Borges PAV, Santos AMC, Elias RB, Gabriel R (2020) The Azores Archipelago: Biodiversity Erosion and Conservation Biogeography. In: Goldstein M, DellaSala D (Eds) Encyclopedia of the World's Biomes. Elsevier. 101-113 pp. <https://doi.org/10.1016/b978-0-12-409548-9.11949-9>
- Borges PAV, Nunes R, Lamelas-López L, Pereira E, Costa R, Monjardino P, Lopes D, Soares AO, Gil A, Rigal F, Ferrante M, Lóvei G (2021) Monitoring Arthropods in Azorean Agroecosystems: the project AGRO-ECOSERVICES. Biodiversity Data Journal 9: e77548. <https://doi.org/10.3897/bdj.9.e77548>
- Borges PAV, Lamelas-Lopez L, Andrade R, Lhoumeau S, Vieira V, Soares A, Borges I, Boieiro M, Cardoso P, Crespo LC, Karsholt O, Schülke M, Serrano ARM, Quartau JA, Assing V (2022) An updated checklist of Azorean arthropods (Arthropoda). Biodiversity Data Journal 10: e97682. <https://doi.org/10.3897/bdj.10.e97682>
- deCastro-Arrazola I, Hortal J, Noriega JA, Sánchez-Piñero F (2020) Assessing the functional relationship between dung beetle traits and dung removal, burial, and seedling emergence. Ecology 101 (10): e03138. <https://doi.org/10.1002/ecy.3138>
- deCastro-Arrazola I, Andrew N, Berg M, Curtsdotter A, Lumaret J, Menéndez R, Moretti M, Nervo B, Nichols E, Sánchez-Piñero F, Santos AC, Sheldon K, Slade E, Hortal J (2023) A trait-based framework for dung beetle functional ecology. Journal of Animal Ecology 92 (1): 44-65. <https://doi.org/10.1111/1365-2656.13829>
- Drăghici A, Ruicănescu A, Cuzepan-Bebeșelea G, Chimișliu C, Mancu C (2019) Distribution of *Onthophagus vacca* and *O. medius* (Coleoptera: Scarabaeidae, Scarabaeinae) in Romania. Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa" 62 (1): 31-41. <https://doi.org/10.3897/travaux.62.e35197>
- Dueñas-Rojas A, Cuesta E, Parmentier LM, Leite A, Borges PAV, Santos AMC (2025) Dung beetles of Azorean cattle-grazed pasturelands – data of the DUNGPOOL project. 1.3. GBIF. Release date: 2025-6-23. URL: https://ipt.gbif.es/resource?r=dungpool_azores
- Fernández-Palacios JM, Kreft H, Irl SDH, Norder S, Ah-Peng C, Borges PAV, Burns KC, de Nascimento L, Meyer J, Montes E, Drake DR (2021) Scientists' warning – The outstanding biodiversity of islands is in peril. Global Ecology and Conservation 31: e01847. <https://doi.org/10.1016/j.gecco.2021.e01847>
- Kadiri N, Lumaret JP, Martínez MI (2022) Funciones ecológicas y servicios ecosistémicos brindados por los escarabajos del estiércol. In: Martínez MI, Lumaret JP (Eds) Escarabajos estercoleros. Biología reproductiva y su regulación. Asociación Española de Entomología. 313-371 pp.
- Lumaret JP, Kadiri N, Martínez MI (2022) Los escarabajos estercoleros y las alteraciones ambientales: riesgos, amenazas y resiliencia de los ecosistemas. In: Martínez MI, Lumaret JP (Eds) Escarabajos del estiércol. Biología reproductiva y su regulación. Asociación Española de Entomología. León, 373-404 pp.
- Mediavilla AC (2024) Seasonal diversity patterns of dung associated beetles on two cow dung insect communities in Terceira Island (Açores, Portugal) [PhD Thesis]. Universidad Barcelona, Barcelona.

- Nichols E, Spector S, Louzada J, Larsen T, Amezcuita S, Favila ME (2008) Ecological functions and ecosystem services provided by Scarabaeinae dung beetles. *Biological Conservation* 141 (6): 1461-1474. <https://doi.org/10.1016/j.biocon.2008.04.011>
- Noriega JA, March-Salas M, Castillo S, García-Q H, Hortal J, Santos AMC (2021) Human perturbations reduce dung beetle diversity and dung removal ecosystem function. *Biotropica* 53 (3): 753-766. <https://doi.org/10.1111/btp.12953>
- Noriega JA, Hortal J, deCastro-Arrazola I, Alves-Martins F, Ortega JG, Bini LM, Andrew N, Arellano L, Beynon S, Davis AV, Favila M, Floate K, Horgan F, Menéndez R, Milotic T, Nervo B, Palestrini C, Rolando A, Scholtz C, Senyüz Y, Wassmer T, Adam R, Araújo C, Barragan-Ramírez JL, Boros G, Camero-Rubio E, Cruz M, Cuesta E, Damborsky MP, Deschodt C, Rajan PD, D'hondt B, Díaz Rojas A, Dindar K, Escobar F, Espinoza V, Ferrer-Paris JR, Gutiérrez Rojas PE, Hemmings Z, Hernández B, Hill S, Hoffmann M, Jay-Robert P, Lewis K, Lewis M, Lozano C, Marín-Armijos D, de Farias PM, Murcia-Ordoñez B, Karimbunkara SN, Navarrete-Heredia JL, Ortega-Echeverría C, Pablo-Cea J, Perrin W, Pessoa MB, Radhakrishnan A, Rahimi I, Raimundo AT, Ramos DC, Rebolledo R, Roggero A, Sánchez-Mercado A, Somay L, Stadler J, Tahmasebi P, Triana Céspedes JD, Santos AMC (2023) Dung removal increases under higher dung beetle functional diversity regardless of grazing intensification. *Nature Communications* 14 (1): 8070. <https://doi.org/10.1038/s41467-023-43760-8>
- Rössner E, Schönfeld J, Ahrens D (2010) *Onthophagus* (Palaeonthophagus) *medius* (Kugelann, 1792)—a good western palaeartic species in the *Onthophagus vacca* complex (Coleoptera: Scarabaeidae: Scarabaeinae: Onthophagini). *Zootaxa* 2629 (1): 1-28. <https://doi.org/10.11646/zootaxa.2629.1.1>
- Roy L, Bon M, Cesarini C, Serin J, Bonato O (2016) Pinpointing the level of isolation between two cryptic species sharing the same microhabitat: a case study with a scarabaeid species complex. *Zoologica Scripta* 45 (4): 407-420. <https://doi.org/10.1111/zsc.12166>
- Tonelli M, Lobo JM (2025) Human activities and dispersal capacity: key drivers of dung beetle colonization in Mediterranean and Macaronesian islands. *Biological Journal of the Linnean Society* 144: blaf003. <https://doi.org/10.1093/biolinnear/blaf003>
- Verdú J, Lobo J, Sánchez-Piñero F, Gallego B, Numa C, Lumaret J, Cortez V, Ortiz A, Tonelli M, García-Teba J, Rey A, Rodríguez A, Durán J (2018) Ivermectin residues disrupt dung beetle diversity, soil properties and ecosystem functioning: An interdisciplinary field study. *Science of The Total Environment* 618: 219-228. <https://doi.org/10.1016/j.scitotenv.2017.10.331>